

STI Indicators by Patient Group

Table of Contents

pg. 2-6 – All Patients

pg. 7 – Men who have sex with men (MSM) Patients

pg. 8 – Transgender Patients

pg. 9 – Women/Pregnant Women

All Patients

Sexual History

Comprehensive Sexual History Elements

Percentage of patients who had a comprehensive sexual history taken.

Denominator	Numerator
All patients.	Number who had a sexual history taken during the review period that includes the elements listed below.

Exclusion(s): None.

- A. Sexual activity during the review period
- B. Number of sexual partners during the review period
- C. Sexual orientation and gender identification
- D. History of STIs in the previous year

If not clearly monogamous:

- E. Sites of sexual exposure (genital, anal, oral)
 - i. For MSM and transgender patients: history of unprotected anal intercourse or other high-risk behaviors
- F. History of condom use/barrier protection

Screening

Syphilis

Syphilis Screening of Sexually Active Patients

Percentage of sexually active patients who received syphilis testing.

Denominator	Numerator
Number of sexually active, non-monogamous patients.	Number who received syphilis testing during the review period.

Exclusion(s): None.

Gonorrhea

Gonorrhea Screening of Sexually Active Patients

Percentage of sexually active patients who had a test for gonorrhea.

Denominator	Numerator
Number of sexually active, non-monogamous patients.	Number who received a gonorrhea test during the review period.

Exclusion(s): None.

Chlamydia

Chlamydia Screening of Sexually Active Patients

Percentage of sexually active patients who received Chlamydia testing.

Denominator	Numerator
Number of sexually active, non-monogamous patients.	Number who received a Chlamydia test during the review period.

Exclusion(s): None.

Genital Chlamydia Testing of Sexual Partners of Individuals with Chlamydia

Percentage of sexual partners of individuals with Chlamydia who had a genital test for Chlamydia.

Denominator	Numerator
Number of sexual partners of individuals with Chlamydia.	Number who had a genital (urine, cervical, urethral, or vaginal) test for Chlamydia performed during the review period.

Exclusion(s): None.

Treatment

Syphilis

Appropriate Treatment of Primary Syphilis

Percentage of patients who received appropriate treatment for primary syphilis following a positive test.

Denominator	Numerator
Number of patients who received treatment following a positive syphilis test.	Number who received a single intramuscular injection of 2.4 million units Benzathine penicillin G during the review period.

Exclusion(s): None.

Gonorrhea

Appropriate Treatment of Gonorrhea

Percentage of patients who received appropriate treatment for gonorrhea following a positive test.

Denominator	Numerator
Number of patients who received treatment following a positive gonorrhea test.	Number who received dual therapy of either 250mg Ceftriaxone and 1g Azithromycin OR 400mg Cefixime and 1g Azithromycin during the review period.

Exclusion(s): None.

Chlamydia

Appropriate Treatment of Chlamydia

Percentage of patients who received appropriate treatment for Chlamydia following a positive test.

Denominator	Numerator
Number of patients who received treatment following a positive Chlamydia test.	Number who received any of the following as treatment for Chlamydia during the review period: <ul style="list-style-type: none">• Single dose of 1g Azithromycin;• 100mg of Doxycycline, 2x daily for 7 days;• 500mg of Erythromycin, 4x daily for 7 days;• 800mg Erythromycin ethylsuccinate, 4x daily for 7 days;• 500mg Levofloxacin, 1x daily for 7 days;• 300mg Ofloxacin, 2x daily for 7 days.

Exclusion(s): If patient is pregnant.

Partner Treatment

Syphilis

Percentage of patients diagnosed with primary syphilis who have had any sexual partner treated for syphilis.

Denominator	Numerator
Number of patients diagnosed with primary syphilis.	Number whose partners have been treated during the review period.

Exclusion(s): None

Percentage of sexual partners of a patient with primary syphilis within 90 days of diagnosis who have been treated.

Denominator	Numerator
Number of sexual partners of a patient with primary syphilis within 90 days of diagnosis.	Number have been treated during the review period.

Exclusion(s): None

Gonorrhea

Percentage of patients diagnosed with gonorrhea who have had any sexual partner treated for gonorrhea.

Denominator	Numerator
Number of patients diagnosed with gonorrhea.	Number whose partners have been treated during the review period.

Exclusion(s): None

Percentage of sexual partners of a patient with gonorrhea within 60 days of diagnosis who have been treated.

Denominator	Numerator
Number of sexual partners of a patient with gonorrhea within 60 days of diagnosis.	Number have been treated during the review period.

Exclusion(s): None

Chlamydia

Percentage of patients diagnosed with chlamydia who have had any sexual partner treated for Chlamydia.

Denominator	Numerator
Number of patients diagnosed with Chlamydia.	Number whose partners have been treated during the review period.

Exclusion(s): None

Percentage of sexual partners of a patient with Chlamydia within 60 days of diagnosis who have been treated.

Denominator	Numerator
Number of sexual partners of a patient with Chlamydia within 60 days of diagnosis.	Number have been treated during the review period.

Exclusion(s): None

Retesting Following Treatment

Syphilis

Percentage of patients who were retested for primary syphilis following treatment.

Denominator	Numerator
Number of patients who received treatment for primary syphilis.	Number of patients who were retested for syphilis at 6 months \pm 14 days and 12 months \pm 14 days following treatment during the review period.

Exclusion(s): None.

Gonorrhea

Percentage of patients who were retested for gonorrhea following treatment.

Denominator	Numerator
Number of patients who received treatment for gonorrhea.	Number of patients who were retested for gonorrhea within 4 months following treatment during the review period.

Exclusion(s): None.

Chlamydia

Percentage of patients who were retested for Chlamydia following treatment.

Denominator	Numerator
Number of patients who received treatment for Chlamydia.	Number of patients who were retested for Chlamydia within 4 months following treatment during the review period.

Exclusion(s): None.

HIV

Known HIV Status or HIV Testing

Percentage of patients who knew their HIV status or received HIV testing.

Denominator	Numerator
All patients aged 13+.	Number who knew their HIV status or had an HIV test performed during the review period.

Exclusion(s): Individual is being treated for a life threatening emergency or the individual lacks capacity to consent to an HIV-related test.

Offering PrEP to At-Risk HIV- Patients

Percentage of at-risk HIV- patients who were offered PrEP.

Denominator	Numerator
Number of at-risk HIV- patients, defined by the table below.	Number who were offered PrEP during the review period.

Exclusion(s): None.

TABLE 1. POTENTIAL CANDIDATES FOR PREP
<p>Clinicians should discuss PrEP with the following non-HIV-infected individuals who have substantial and ongoing risk:</p> <ul style="list-style-type: none">• Men who have sex with men (MSM) who engage in unprotected anal intercourse^{15,16}• Individuals who are in a serodiscordant sexual relationship with a known HIV-infected partner• Male-to-female and female-to male transgender individuals engaging in high-risk sexual behaviors• Individuals engaging in transactional sex, such as sex for money, drugs, or housing• Injection drug users who report any of the following behaviors: sharing injection equipment (including to inject hormones among transgender individuals), injecting one or more times per day, injecting cocaine or methamphetamine, engaging in high-risk sexual behaviors¹³• Individuals who use stimulant drugs associated with high-risk behaviors, such as methamphetamine¹⁵⁻¹⁸• Individuals diagnosed with at least one anogenital sexually transmitted infection in the last year^{19,20}• Individuals who have been prescribed non-occupational post-exposure prophylaxis (nPEP) who demonstrate continued high-risk behavior or have used multiple courses of nPEP²¹

Other individuals may qualify for PrEP who may not fit within the above risk categories. Decisions to initiate PrEP should be individualized by weighing patients' personal risk of acquiring HIV infection against the potential benefits and risks of TDF/FTC.

Sources Used:

[A Guide to Taking a Sexual History](#) – CDC Publication

[2015 Sexually Transmitted Diseases Treatment Guidelines](#) – CDC

MSM Patients

Semiannual Syphilis Testing of MSM

Percentage of MSM who received semiannual testing for syphilis.

Denominator	Numerator
Number of MSM patients.	Number who had at least one test for syphilis performed during each 6-month period of the review period.

Exclusion(s): MSM patients only seen annually.

Annual Extragenital Gonorrhea Testing Among MSM and MtF Transgender Patients

Percentage of MSM or MtF transgender patients who receive annual extragenital testing for gonorrhea.

Denominator	Numerator
Number of MSM or MtF transgender patients.	Number who had an annual extragenital (anorectal and pharyngeal) test for gonorrhea performed during the review period.

Exclusion(s): None.

Annual Genital Chlamydia Testing Among MSM, MtF Transgender Patients, and Women

Percentage of MSM or MtF transgender patients or women who have an annual genital test for Chlamydia.

Denominator	Numerator
Number of MSM or MtF transgender patients or female patients.	Number who had an annual genital (urine, cervical, urethral, or vaginal) test for Chlamydia performed during the review period.

Exclusion(s): None.

Annual Extragenital Chlamydia Testing Among MSM and MtF Transgender Patients

Percentage of MSM or MtF transgender patients who receive annual extragenital testing for Chlamydia.

Denominator	Numerator
Number of MSM or MtF transgender patients.	Number who had an annual extragenital (anorectal and pharyngeal) test for Chlamydia performed during the review period.

Exclusion(s): None.

Transgender Patients

Testing

Annual Extragenital Gonorrhea Testing Among MSM and MtF Transgender Patients

Percentage of MSM or MtF transgender patients who receive annual extragenital testing for gonorrhea.

Denominator	Numerator
Number of MSM or MtF transgender patients.	Number who had an annual extragenital (anorectal and pharyngeal) test for gonorrhea performed during the review period.

Exclusion(s): None.

Annual Genital Chlamydia Testing Among MSM, MtF Transgender Patients, and Women

Percentage of MSM or MtF transgender patients or women who have an annual genital test for Chlamydia.

Denominator	Numerator
Number of MSM or MtF transgender patients or female patients.	Number who had an annual genital (urine, cervical, urethral, or vaginal) test for Chlamydia performed during the review period.

Exclusion(s): None.

Annual Extragenital Chlamydia Testing Among MSM and MtF Transgender Patients

Percentage of MSM or MtF transgender patients who receive annual extragenital testing for Chlamydia.

Denominator	Numerator
Number of MSM or MtF transgender patients.	Number who had an annual extragenital (anorectal and pharyngeal) test for Chlamydia performed during the review period.

Exclusion(s): None.

Women/Pregnant Women

Testing

Syphilis Testing of Pregnant Women

Percentage of pregnant female patients who had a test for syphilis.

Denominator	Numerator
Number of pregnant female patients.	Number who had a test for syphilis during the review period.

Exclusion(s): None.

Annual Genital Chlamydia Testing Among MSM, MtF Transgender Patients, and Women

Percentage of MSM or MtF transgender patients or women who have an annual genital test for Chlamydia.

Denominator	Numerator
Number of MSM or MtF transgender patients or female patients.	Number who had an annual genital (urine, cervical, urethral, or vaginal) test for Chlamydia performed during the review period.

Exclusion(s): None.

Genital Chlamydia Testing Among Pregnant Women

Percentage of pregnant women who had genital tests for Chlamydia during the 1st and 3rd trimesters of pregnancy.

Denominator	Numerator
Number of pregnant female patients.	Number who had genital (urine, cervical, urethral, or vaginal) tests for Chlamydia performed during the 1 st and 3 rd trimesters of pregnancy during the review period.

Exclusion(s): None.

Treatment

Appropriate Treatment of Chlamydia – Pregnant Women

Percentage of pregnant women who received appropriate treatment for Chlamydia following a positive test.

Denominator	Numerator
Number of pregnant patients who received treatment following a positive Chlamydia test.	Number who received any of the following as treatment for Chlamydia during the review period: <ul style="list-style-type: none">• Single dose of 1g Azithromycin;• 500mg Amoxicillin, 3x daily for 7 days;• 500mg Erythromycin, 4x daily for 7 days;• 250mg Erythromycin, 4x daily for 14 days;• 800mg Erythromycin ethylsuccinate, 4x daily for 7 days;• 400mg Erythromycin ethylsuccinate, 4x daily for 14 days.

Exclusion(s): None.